

Determination of the Gaugino Mass Parameter M_1 in Different Linear Collider Modes

Claus Blöchinger

in collaboration with

Hans Fraas, Gudrid Moortgat-Pick and Tobias
Mayer

University of Würzburg

LCWS, Fermilab, 25.10.2000

Contents

I. Introduction and Motivation

II. Some References

III. M_1 Dependence of SUSY Processes

III.1. Neutralino Pair Production in e^+e^-

III.2. LSP/Selectron Production in $e^-\gamma$

III.3. Chargino Pair Production in $\gamma\gamma$

III.4. Selectron Pair Production in e^-e^-

IV. Conclusions and Outlook

I. Introduction and Motivation

First detection of SUSY:

L(arge) H(adron) C(ollider): $\tilde{q}, \tilde{g} < 2.5$ TeV

What is left for a Linear Collider:

precise determination of underlying SUSY parameters

Unconstrained SUSY: **105** free parameters !

- to reduce parameters: assumptions on their behavior at the GUT-scale, CP, FCNC,...
- few free parameters, e.g. gaugino masses M_1 , M_2 and M_3

One usual Assumption:

Unification of M_1 , M_2 and M_3 at the GUT-Scale



RG equations



$$M_1 = \frac{5}{3} \tan^2 \theta_W M_2$$

But:

- Unification - assumption must not be right!
- Different SUSY breaking scenarios:

F_Φ	M_1	M_2	M_3
1	1(~ 6)	1(~ 2)	1(~ 1)
24	2(~ 12)	-3(~ -6)	-1(~ -1)
75	1(~ 6)	3(~ 6)	-5(~ -5)
200	1(~ 6)	2(~ 4)	10(~ 10)
OII	1(~ 6)	5(~ 10)	53/5($\sim 53/5$)

Chen, Drees, Gunion, 1997/1999

→ M_1 determination necessary

Particle Content of the MSSM

Particles of the SM	SUSY Particles	
	Int-ES	Mass-ES
$l = e, \mu, \tau$	\tilde{l}_L, \tilde{l}_R	\tilde{l}_1, \tilde{l}_2 Slepton
$\nu = \nu_e, \nu_\mu, \nu_\tau$	$\tilde{\nu}$	$\tilde{\nu}$ Sneutrino
$q = u, d, s, c, b, t$	\tilde{q}_L, \tilde{q}_R	\tilde{q}_1, \tilde{q}_2 Squark
W^\pm	\tilde{W}^\pm	$\tilde{\chi}_j^\pm$ Chargino
H^\pm	\tilde{H}^\pm	
γ	$\tilde{\gamma}$	
Z	\tilde{Z}	$\tilde{\chi}_j^0$ Neutralino
H_1^0	\tilde{H}_1^0	
H_2^0	\tilde{H}_2^0	
g	\tilde{g}	Gluino

$$\tilde{\chi}_i^0 = N_{i1}\tilde{\gamma} + N_{i2}\tilde{Z} + N_{i3}\tilde{H}_1^0 + N_{i4}\tilde{H}_2^0$$

Parameters of the MSSM

- M_2 : SU(2) gaugino mass parameter
- M_1 : U(1) gaugino mass parameter
- μ : Higgsino mass parameter
- $\tan \beta = \frac{v_2}{v_1}$: ratio of the Higgs vacuum expectation values
- $m_{\tilde{f}_{L/R}}$: sfermion masses (corr. to m_0 , the common sfermion mass at the GUT scale)

"Usual" GUT-Relation:

$$M_1 = \frac{5}{3} \tan^2 \theta_W M_2$$

Neutralino - Mixing

Mass matrix of the neutralinos:

$$\mathcal{Y} = \begin{pmatrix} M_2 \sin^2 \theta_W + \textcolor{red}{M}_1 \cos^2 \theta_W & (M_2 - \textcolor{red}{M}_1) \sin \theta_W \cos \theta_W & 0 & 0 \\ (M_2 - \textcolor{red}{M}_1) \sin \theta_W \cos \theta_W & M_2 \cos^2 \theta_W + \textcolor{red}{M}_1 \sin^2 \theta_W & m_Z & 0 \\ 0 & m_Z & \mu \sin 2\beta & -\mu \cos 2\beta \\ 0 & 0 & -\mu \cos 2\beta & -\mu \sin 2\beta \end{pmatrix}$$

Masses of the neutralinos:


$$\begin{aligned} & \left(m_{\tilde{\chi}_i^0}^2 - \mu^2 \right) \left(m_{\tilde{\chi}_i^0}^2 - M_2 \right) \left(m_{\tilde{\chi}_i^0}^2 - \textcolor{red}{M}_1 \right) - m_Z^2 \left(m_{\tilde{\chi}_i^0}^2 + \mu \sin 2\beta \right) \cdot \\ & \cdot \left(m_{\tilde{\chi}_i^0}^2 - M_2 \sin^2 \theta_W - \textcolor{red}{M}_1 \cos^2 \theta_W \right) = 0 \end{aligned}$$

Neutralino Eigenstates:

$$\tilde{\chi}_i^0 = \frac{1}{N_i} \left(\begin{array}{l} (M_2 - \textcolor{red}{M}_1) \sin \theta_W \cos \theta_W \left(m_{\tilde{\chi}_i^0}^2 - \mu^2 \right) \\ \left(m_{\tilde{\chi}_i^0}^2 - M_2 \sin^2 \theta_W - \textcolor{red}{M}_1 \cos^2 \theta_W \right) \left(m_{\tilde{\chi}_i^0}^2 - \mu^2 \right) \\ \left(m_{\tilde{\chi}_i^0}^2 - M_2 \sin^2 \theta_W - \textcolor{red}{M}_1 \cos^2 \theta_W \right) \left(m_{\tilde{\chi}_i^0}^2 + \mu \sin 2\beta \right) \\ - \left(m_{\tilde{\chi}_i^0}^2 - M_2 \sin^2 \theta_W - \textcolor{red}{M}_1 \cos^2 \theta_W \right) \mu \cos 2\beta \end{array} \right)$$

$$N_i = \{ \left(m_{\tilde{\chi}_i^0}^2 - \mu^2 \right)^2 \left(\sin^2 \theta_W \left(m_{\tilde{\chi}_i^0}^2 - M_2 \right)^2 + \cos^2 \theta_W \left(m_{\tilde{\chi}_i^0}^2 - \textcolor{red}{M}_1 \right)^2 \right) + \\ + \left(m_{\tilde{\chi}_i^0}^2 - M_2 \right)^2 \left(m_{\tilde{\chi}_i^0}^2 - \textcolor{red}{M}_1 \right)^2 m_Z^{-2} \} + \\ + \left(m_{\tilde{\chi}_i^0}^2 - M_2 \sin^2 \theta_W - \textcolor{red}{M}_1 \cos^2 \theta_W \right)^2 (m_Z \mu \cos 2\beta)^2 \}^{\frac{1}{2}}$$

Neutralino Masses $m_{\tilde{\chi}_i^0}$



$$M_2 = 152 \text{ GeV}$$

$$\mu = 316 \text{ GeV}$$

$$\tan \beta = 3$$


II. Some References

- MSSM without M_1/M_2 GUT-Relation
C.H. Chen, M. Drees, J.F. Gunion, Phys.Rev.Lett. 76 (1996) 2002
C.H. Chen, M. Drees, J.F. Gunion, Phys.Rev. D55 (1997) 330;
Erratum-ibid. D60 (1999) 039901
Snowmass (1996)
- Monte Carlo Simulations
J.L. Feng, M.J. Strassler, Phys.Rev. D51 (1995) 4661
J.L. Feng, M.J. Strassler, Phys.Rev. D55 (1997) 1326
T. Tsukamoto, et al., Phys.Rev. D51 (1995) 3153
M.M. Nojiri, K. Fujii, T. Tsukamoto, Phys.Rev. D54 (1996) 6756
- e^-e^- Collisions
M.E. Peskin, Int.J.Mod.Phys. A13 (1998) 2299-2306
J.L. Feng, Int.J.Mod.Phys. A13 (1998) 2319
- “Analytical” from Chargino/Neutralino Masses
J.-L. Kneur, G. Moultaka, Phys.Rev. D59 (1999) 015005
- “Analytical” from Chargino Production and $m_{\tilde{\chi}_1^0}$
S.Y. Choi, et al, Eur.Phys.J. C7 (1999) 123-134
S.Y. Choi, et al, Eur.Phys.J. C8 (1999) 669-677
- Processes in this talk

III. M_1 -dependence of SUSY Processes

MSSM-Parameters:

$$M_1, \quad M_2, \quad \mu, \tan \beta, \quad m_{\tilde{e}_L}, \quad m_{\tilde{e}_R}$$



GUT-Relation: $M_1 = M_2 \cdot \frac{5}{3} \tan^2 \theta_W$

CP-Conservation: M_1, M_2, μ real

Parameters for all processes:

$$M_2 = 152 \text{ GeV}$$

$$\mu = 316 \text{ GeV}$$

$$\tan \beta = 3$$

Different choices for:

$\sqrt{s_{ee}}$, $m_{\tilde{e}_L}$, $m_{\tilde{e}_R}$, polarized e^+e^-/γ beam

III.1. Neutralino Production in e^+e^- Annihilation


Process:

$$e^- + e^+ \xrightarrow[\tilde{e}_{L/R}]{Z^0} \tilde{\chi}_1^0 + \tilde{\chi}_2^0$$
$$\quad\quad\quad \downarrow \quad\quad\quad \xrightarrow[\tilde{e}_{L/R}]{Z^0} \tilde{\chi}_1^0 + e^+ + e^-$$


- Associated production of $\tilde{\chi}_1^0$ and $\tilde{\chi}_2^0$ with leptonic decay of $\tilde{\chi}_2^0$
- Complete spin correlations between production and decay:
 - + contributions independent of $\tilde{\chi}_2^0$ -polarisation
 - + contributions depending on $\tilde{\chi}_2^0$ -polarisation

Feynman Graphs for Neutralino Production and Leptonic Decay

Production :



Decay :



M_1 in Production and Decay:

- masses $m_{\tilde{\chi}_1^0}, m_{\tilde{\chi}_2^0}$
- couplings $O''_{12}, f_{e1}^L f_{e2}^L, f_{e1}^R f_{e2}^R$

Couplings for the Different Processes

$$O_{11}^L = -\frac{1}{\sqrt{2}} (\cos \beta N_{14} - \sin \beta N_{13}) V_{12} + (\sin \theta_W N_{11} + N_{12}) V_{11}^*$$

$$O_{11}^R = \frac{1}{\sqrt{2}} (\sin \beta N_{14} - \cos \beta N_{13}^*) U_{12} + (\sin \theta_W N_{11}^* + N_{12}^*) U_{11}$$

$$O_{12}''^L = -\frac{1}{2} (N_{13} N_{23}^* - N_{14} N_{24}^*) \cos 2\beta - \frac{1}{2} (N_{13} N_{24}^* + N_{14} N_{23}^*) \sin 2\beta$$

$$O_{12}''^R = -O_{12}''^{L*}$$


$$f_{ei}^L = -\sqrt{2} \left(\frac{1}{\cos \theta_W} (T_{3e} + \sin^2 \theta_W) N_{i2} - \sin \theta_W N_{i1} \right)$$

$$f_{ei}^R = \sqrt{2} \sin \theta_W (\tan \theta_W N_{i2}^* - N_{i1}^*)$$

$$f_{\nu 1}^L = -\frac{\sqrt{2}}{\cos \theta_W} T_{3\nu} N_{12}$$

Total Cross Section σ_e

$$\sigma_e = \sigma(e^+e^- \rightarrow \tilde{\chi}_1^0\tilde{\chi}_2^0) \times BR(\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 e^+e^-)$$



$$\sqrt{s_{ee}} = m_{\tilde{\chi}_1^0} + m_{\tilde{\chi}_2^0} + 30 \text{ GeV}$$

$$m_{\tilde{e}_R} = 161 \text{ GeV}$$

$$m_{\tilde{e}_L} = 176 \text{ GeV}$$


$$P_- = 0, \pm 0.85, P_+ = 0, \pm 0.6 \rightarrow (\text{sgn}(P_-), \text{sgn}(P_+))$$

Asymmetry A_{FB} of the decay electron

$$A_{FB} = \frac{\sigma_e (\cos \Theta_e > 0) - \sigma_e (\cos \Theta_e < 0)}{\sigma_e (\cos \Theta_e > 0) + \sigma_e (\cos \Theta_e < 0)}$$

A_{FB}

G. Moortgat-Pick, 1999



$$\sqrt{s_{ee}} = m_{\tilde{\chi}_1^0} + m_{\tilde{\chi}_2^0} + 30 \text{ GeV}$$


$$m_{\tilde{e}_R} = 161 \text{ GeV}$$

$$m_{\tilde{e}_L} = 176 \text{ GeV}$$

$$P_- = 0, \pm 0.85, P_+ = 0, \pm 0.6 \rightarrow (\text{sgn}(P_-), \text{sgn}(P_+))$$

Polarization Asymmetry A_{pol}

$$A_{pol} = \frac{\sigma_{ee}(+0) - \sigma_{ee}(-0)}{\sigma_{ee}(+0) + \sigma_{ee}(-0)}$$



$$\sqrt{s_{ee}} = m_{\tilde{\chi}_1^0} + m_{\tilde{\chi}_2^0} + 50 \text{ GeV}$$

$$m_{\tilde{e}_R} = 161 \text{ GeV}$$


$$m_{\tilde{e}_L} = 176 \text{ GeV}$$

$$P_- = \pm 0.9, P_+ = 0$$

III.2. LSP/Selectron Production

in $e\gamma$ Scattering

Process:



- LSP/Selectron production with leptonic decay
- Photon beam: Compton-backscattering of laser photons
- Higher mass reach in $m_{\tilde{e}_{L/R}}$ for $e\gamma$ -mode
- For the plots in this section:
 - P_e = long. polarization of the electron beam
 - λ_L = circular polarization of the laser beam
 - λ_k = long. polarization of the converted electron beam

Observable σ_{ee} :

Total cross section in the $e\gamma$ -cms:

$$\sigma_{e\gamma}^{L/R} = \sigma_P(s_{e\gamma}) \cdot BR(\tilde{e}_{L/R} \rightarrow e^- \tilde{\chi}_1^0)$$

Important relation:

$$\boxed{\sigma_{e\gamma}^{L/R} \propto \left(f_1^{L/R}\right)^2}$$

Convolution

Total cross section in the ee -cms:

$$\sigma_{ee}^{L/R} = \int dy P_C(y) \hat{\sigma}_{e\gamma}^{L/R}(s_{e\gamma} = ys_{ee})$$

$P_C(y = \frac{E_\gamma}{E_k})$ = γ - spectrum of the Compton backscattered laser photons

→ λ_k = helicity of the converted e^-

→ λ_L = helicity of the laser photons

$$\hat{\sigma}_{e\gamma}^{L/R} = \frac{1}{2}(1 + \lambda(y)) \left(\sigma_{e\gamma}^{L/R}\right)^+ + \frac{1}{2}(1 - \lambda(y)) \left(\sigma_{e\gamma}^{L/R}\right)^-$$

$\left(\sigma_{e\gamma}^{L/R}\right)^{+/-}$ = total cross sections for
a completely right (left)
circular polarized photon beam

$\lambda(y)$ = mean helicity of the photon beam

Same final states for \tilde{e}_R and \tilde{e}_L

$$\rightarrow \sigma_{ee} = \sigma_{ee}^L + \sigma_{ee}^R$$

Polarisation Asymmetries:


$$A_e = \frac{\sigma_{ee}(+P_e) - \sigma_{ee}(-P_e)}{\sigma_{ee}(+P_e) + \sigma_{ee}(-P_e)}$$

$$A_{\lambda_L} = \frac{\sigma_{ee}(+\lambda_L) - \sigma_{ee}(-\lambda_L)}{\sigma_{ee}(+\lambda_L) + \sigma_{ee}(-\lambda_L)}$$


$$A_\gamma = \frac{\sigma_{ee}(+\lambda_L, -\lambda_k) - \sigma_{ee}(-\lambda_L, +\lambda_k)}{\sigma_{ee}(+\lambda_L, -\lambda_k) + \sigma_{ee}(-\lambda_L, +\lambda_k)}$$

Feynman Graphs for LSP/Selectron Production and Leptonic Decay

Production :



Decay :




M_1 in Production and Decay:

- mass $m_{\tilde{\chi}_1^0}$
- couplings f_{e1}^L , f_{e1}^R

Polarization Asymmetry A_γ

$$A_\gamma = \frac{\sigma_{ee}(s_{ee}, P_e, -\lambda_k, +\lambda_L) - \sigma_{ee}(s_{ee}, P_e, +\lambda_k, -\lambda_L)}{\sigma_{ee}(s_{ee}, P_e, -\lambda_k, +\lambda_L) + \sigma_{ee}(s_{ee}, P_e, +\lambda_k, -\lambda_L)}$$



$$\sqrt{s_{ee}} = 500 \text{ GeV}$$

$$|\lambda_k| = 0.8$$

$$|\lambda_L| = 1$$


$$P_e = 0.8$$

$$m_{\tilde{e}_L} = 179.3 \text{ GeV}$$

$$m_{\tilde{e}_R} = 137.7 \text{ GeV}$$

Polarization Asymmetry A_{λ_L}

$$A_{\lambda_L} = \frac{\sigma_{ee}(s_{ee}, P_e, \lambda_k, \lambda_L) - \sigma_{ee}(s_{ee}, P_e, \lambda_k, -\lambda_L)}{\sigma_{ee}(s_{ee}, P_e, \lambda_k, \lambda_L) + \sigma_{ee}(s_{ee}, P_e, \lambda_k, -\lambda_L)}$$



$$\sqrt{s_{ee}} = 500 \text{ GeV}$$

$$\lambda_L = +1$$

$$\lambda_k = +0.8$$


$$P_e = 0.8$$

$$m_{\tilde{e}_L} = 179.3 \text{ GeV}$$

$$m_{\tilde{e}_R} = 137.7 \text{ GeV}$$

Polarization Asymmetry A_{λ_L}

$$A_{\lambda_L} = \frac{\sigma_{ee}(s_{ee}, P_e, \lambda_k, \lambda_L) - \sigma_{ee}(s_{ee}, P_e, \lambda_k, -\lambda_L)}{\sigma_{ee}(s_{ee}, P_e, \lambda_k, \lambda_L) + \sigma_{ee}(s_{ee}, P_e, \lambda_k, -\lambda_L)}$$



$$\sqrt{s_{ee}} = 500 \text{ GeV}$$

$$\lambda_L = +1$$

$$\lambda_k = -0.8$$

$$P_e = -0.8$$

$$m_{\tilde{e}_L} = 179.3 \text{ GeV}$$

$$m_{\tilde{e}_R} = 137.7 \text{ GeV}$$


Total Cross Section

$\sigma_{ee} (\lambda_L = \pm 1)$

Convoluted cross section in the lab. frame:

$$\sigma_{ee} (s_{ee}, P_e, \lambda_k, \lambda_L) = \int dy P(y) \sigma_{e\gamma} (s_{e\gamma}) \times BR(\tilde{e}_{L/R} \rightarrow \tilde{\chi}_1^0 e^-)$$

σ_{ee}/fb



$$\sqrt{s_{ee}} = 500 \text{ GeV}$$

$$P_e = 0.8$$

$$\lambda_k = +0.8$$

$$\lambda_L = +1 \text{ ---}$$

$$\lambda_L = -1 \text{ - - -}$$

$$m_{\tilde{e}_L} = 179.3 \text{ GeV}$$


$$m_{\tilde{e}_R} = 137.7 \text{ GeV}$$

Total Cross Section

$\sigma_{ee} (\lambda_L = \pm 1)$

Convoluted cross section in the lab. frame:

$$\sigma_{ee} (s_{ee}, P_e, \lambda_k, \lambda_L) = \int dy P(y) \sigma_{e\gamma} (s_{e\gamma}) \times BR(\tilde{e}_{L/R} \rightarrow \tilde{\chi}_1^0 e^-)$$



$$\sqrt{s_{ee}} = 500 \text{ GeV}$$

$$P_e = -0.8$$

$$\lambda_k = -0.8$$

$$\lambda_L = +1 \text{ ---}$$

$$\lambda_L = -1 \text{ - - -}$$

$$m_{\tilde{e}_L} = 179.3 \text{ GeV}$$

$$m_{\tilde{e}_R} = 137.7 \text{ GeV}$$


Total Cross Section

$$\sigma_{ee} (\lambda_L = \pm 1)$$

Convoluted cross section in the lab. frame:

$$\sigma_{ee} (s_{ee}, P_e, \lambda_k, \lambda_L) = \int dy P(y) \sigma_{e\gamma} (s_{e\gamma}) \times BR(\tilde{e}_{L/R} \rightarrow \tilde{\chi}_1^0 e^-)$$

σ_{ee}/fb



$$\sqrt{s_{ee}} = 500 \text{ GeV}$$

$$\rightarrow \sqrt{s_{e\gamma}}^{max} = 0.91 \sqrt{s_{ee}} = 455.1 \text{ GeV}$$

$$m_{\tilde{e}_L} = 350.0 \text{ GeV}, m_{\tilde{e}_R} = 330.5 \text{ GeV}$$


$$\rightarrow m_{\tilde{\chi}_1^0} = 124.6 \text{ GeV} \rightarrow M_1^{max} = 184 \text{ GeV}$$

$$P_e = 0.8 \quad \lambda_k = +0.8$$

$$\lambda_L = +1 \text{ ---} \quad \lambda_L = -1 \text{ - - - -}$$

III.3. Chargino Pair Production in $\gamma\gamma$ Scattering

Process:



- Chargino pair production with leptonic decay
- Complete spin correlations between production and decay:
 - + contributions independent of $\tilde{\chi}_1^\pm$ -polarisation
 - + contributions depending on $\tilde{\chi}_1^+/\tilde{\chi}_1^-$ -polarisation
 - + contributions depending on spin correlations between $\tilde{\chi}_1^+$ and $\tilde{\chi}_1^-$

- In the production **only** QED!

- For the plots in this section:

$\lambda_{L_1}, \lambda_{L_2}$ = circular polarization of the laser beams

$\lambda_{k_1}, \lambda_{k_2}$ = longitudinal polarization of the converted electron beams

Observables

Convoluted cross section in the lab. frame:


$$\sigma_{ee} (s_{ee}, , \lambda_{k_{1,2}}, \lambda_{L_{1,2}}) = \int dy_1 dy_2 P(y_1) P(y_2) \sigma_{\gamma\gamma} \times BR(\tilde{\chi}_1^+ \rightarrow \tilde{\chi}_1^0 e^+ \nu)$$

Forward-backward asymmetry:


$$A_{FB} = \frac{\sigma_e (\cos \Theta_e > 0) - \sigma_e (\cos \Theta_e < 0)}{\sigma_e (\cos \Theta_e > 0) + \sigma_e (\cos \Theta_e < 0)}$$

Feynman Graphs for Chargino Pair Production and Leptonic Decay

Production :



Decay :




M_1 only in the Decay:

- mass $m_{\tilde{\chi}_1^0}$
- couplings $O_{11}^{L/R}$, f_{e1}^L , $f_{\nu 1}^L$

Total Cross Section σ_{ee}

Convoluted cross section in the lab. frame:

$$\sigma_{ee}(s_{ee}, \lambda_{k_{1,2}}, \lambda_{L_{1,2}}) = \int dy_1 dy_2 P(y_1) P(y_2) \sigma_{\gamma\gamma} \times BR(\tilde{\chi}_1^+ \rightarrow \tilde{\chi}_1^0 e^+ \nu)$$



$\sqrt{s_{ee}} = 1000 \text{ GeV}$ (\rightarrow higher cross sections than for $\sqrt{s_{ee}} = 500 \text{ GeV}$)

$$m_{\tilde{e}_R} = 137.7 \text{ GeV}$$

$$m_{\tilde{e}_L} = 179.3 \text{ GeV}$$

$$\lambda_{L_1} = \lambda_{L_2} = 0$$

$$\lambda_{k_1} = 0.8$$


$$m_{\tilde{e}_R} = 330.5 \text{ GeV}$$

$$m_{\tilde{e}_L} = 350.0 \text{ GeV}$$

$$\lambda_{k_2} = -0.8$$

Asymmetry A_{FB} of the decay electron

$$A_{FB} = \frac{\sigma_e (\cos \Theta_e > 0) - \sigma_e (\cos \Theta_e < 0)}{\sigma_e (\cos \Theta_e > 0) + \sigma_e (\cos \Theta_e < 0)}$$



$\sqrt{s_{ee}} = 1000 \text{ GeV}$ (\rightarrow smaller A_{FB} than for $\sqrt{s_{ee}} = 500 \text{ GeV}$)

$$m_{\tilde{e}_R} = 137.7 \text{ GeV}$$

$$m_{\tilde{e}_L} = 179.3 \text{ GeV}$$

$$\lambda_{L_1} = \lambda_{L_2} = 0$$

$$\lambda_{k_1} = 0.8$$

$$m_{\tilde{e}_R} = 330.5 \text{ GeV}$$


$$m_{\tilde{e}_L} = 350.0 \text{ GeV}$$

$$\lambda_{k_2} = -0.8$$

III.4. Selectron Pair Production

in e^-e^- Scattering


Process:




- All 4 neutralinos as exchange particles
- Polarized electron beams
→ better polarisation than in e^+e^-
- For the plots in this section:
 $P_{e1,2}$ = long. polarization of the electron beams

Feynman Graphs for Selectron Pair Production and Decay

Production :



Decay :




M_1 in Production and Decay:

- mass $m_{\tilde{\chi}_i^0}$
- couplings f_{ei}^L, f_{ei}^R

Total Cross Section σ_{ee}

$$\sigma_{ee} = \sigma \left(e^- e^- \rightarrow \tilde{e}_{L/R} \tilde{e}_{L/R} \rightarrow e^- e^- \tilde{\chi}_1^0 \tilde{\chi}_1^0 \right)$$



$$\sigma_{ee} = \sigma_{RR} + \sigma_{LL} + \sigma_{LR} + \sigma_{RL}$$

$$\sqrt{s_{ee}} = 500 \text{ GeV}$$


$$m_{\tilde{e}_R} = 137.7 \text{ GeV}$$

$$m_{\tilde{e}_L} = 179.3 \text{ GeV}$$

$$P_{e_1} = P_{e_2} = 0\%$$

Total Cross Section σ_{ee}

$$\sigma_{ee} = \sigma \left(e^- e^- \rightarrow \tilde{e}_{L/R} \tilde{e}_{L/R} \rightarrow e^- e^- \tilde{\chi}_1^0 \tilde{\chi}_1^0 \right)$$



$$\sigma_{ee} = \sigma_{RR} + \sigma_{LL} + \sigma_{LR} + \sigma_{RL}$$

$$\sqrt{s_{ee}} = 500 \text{ GeV}$$


$$m_{\tilde{e}_R} = 137.7 \text{ GeV}$$

$$m_{\tilde{e}_L} = 179.3 \text{ GeV}$$

$$P_{e_1} = P_{e_2} = 80\%$$

Total Cross Section σ_{ee}

$$\sigma_{ee} = \sigma \left(e^- e^- \rightarrow \tilde{e}_{L/R} \tilde{e}_{L/R} \rightarrow e^- e^- \tilde{\chi}_1^0 \tilde{\chi}_1^0 \right)$$



$$\sigma_{ee} = \sigma_{RR} + \sigma_{LL} + \sigma_{LR} + \sigma_{RL}$$

$$\sqrt{s_{ee}} = 1000 \text{ GeV}$$


$$m_{\tilde{e}_R} = 330 \text{ GeV}$$

$$m_{\tilde{e}_L} = 350 \text{ GeV}$$

$$P_{e_1} = P_{e_2} = 0\%$$

Total Cross Section σ_{ee}

$$\sigma_{ee} = \sigma \left(e^- e^- \rightarrow \tilde{e}_{L/R} \tilde{e}_{L/R} \rightarrow e^- e^- \tilde{\chi}_1^0 \tilde{\chi}_1^0 \right)$$



$$\sigma_{ee} = \sigma_{RR} + \sigma_{LL} + \sigma_{LR} + \sigma_{RL}$$

$$\sqrt{s_{ee}} = 1000 \text{ GeV}$$

$$m_{\tilde{e}_R} = 330 \text{ GeV}$$

$$m_{\tilde{e}_L} = 350 \text{ GeV}$$

$$P_{e_1} = P_{e_2} = 80\%$$

IV. Conclusions and Outlook

- Determination of the gaugino mass parameter M_1 in principle possible in every mode of a linear collider
- $e^+e^- \rightarrow \tilde{\chi}_1^0\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0\tilde{\chi}_1^0 e^+e^-$:
 σ_e , A_{FB} sensitively dependent on M_1 .
Complicated M_1 -dependence
→ ambiguities for $40 \text{ GeV} < M_1 < 160 \text{ GeV}$.
Possibility to separate ambiguities via A_{pol} (depends on selectron masses and splittings).
- $e\gamma \rightarrow \tilde{\chi}_1^0\tilde{e} \rightarrow \tilde{\chi}_1^0\tilde{\chi}_1^0 e^-$:
 σ_{ee} , A_{λ_L} sensitively dependent on M_1 for $40 \text{ GeV} < M_1 < 300 \text{ GeV}$.
With both observables: no ambiguities.

- $\gamma\gamma \rightarrow \tilde{\chi}_1^+ \tilde{\chi}_1^- \rightarrow \tilde{\chi}_1^- \tilde{\chi}_1^0 e^+ \nu_e$:
 strong dependence of σ_e , A_{FB} on M_1
 only for $40 \text{ GeV} < M_1 < 160 \text{ GeV}$.
 With both observables: no ambiguities.
- $e^- e^- \rightarrow \tilde{e}\tilde{e} \rightarrow e^- e^- \tilde{\chi}_1^0 \tilde{\chi}_1^0$:
 Widest range for M_1 ($40 \text{ GeV} < M_1 < 1000 \text{ GeV}$)
 Ambiguities, separation via asymmetries?
- Accuracy between $O(0.1\%)$ and $O(10\%)$ depending on the mode and the parameter point
- Polarization is necessary in every mode for M_1 -determination !
- Outlook: selectron pair production in $e^- e^-$
 selectron pair production in $\gamma\gamma$
 ISR